

Seasonal Moth Trapping for Detection of Adult Flights for Southwestern Corn Borer, Western Bean Cutworm, and Fall Armyworm in the Texas High Plains

Final Report

Dr. Ed Bynum

Extension Entomologist
Texas A&M AgriLife Research and Extension Center
6500 Amarillo Blvd., West
Amarillo, TX 79106
(806) 677- 5600 (FAX 806-677-5644)
Ebynum@ag.tamu.edu

Dr. Pat Porter

Extension Entomologist
Texas A&M AgriLife Research and Extension Center
1102 East FM 1294
Lubbock, Texas 79403-6603
(806) 746 – 6101 (FAX 806-746-4057)

Collaborators

Mr. Rick Auckerman, CEA-Ag, Deaf Smith County
Ms. Nancy Anderson, CEA-Ag, Castro County
Mr. Mike Bragg, CEA-Ag, Hartley/Dallam Counties
Mr. Gary Cross, CEA-Ag, Hale County
Mr. Marcel Fischbacher, CEA-Ag, Moore County
Mr. John Villalba, CEA-Ag, Swisher County
Mr. Brandon McGinty, CEA-Ag, Gray County
Mr. J. D. Ragland, CEA-Ag, Randall County
Mr. J. R. Sprague, CEA-Ag, Lipscomb County
Mr. Scott Strawn, CEA-Ag, Ochiltree County
Ms. Kristy Slough, Hutchinson County

Summary

The Moth Trapping project has been conducted during the 2011 – 2014 growing seasons from June through August to monitor the seasonal moth flights of Southwestern corn borer (SWCB), Western bean cutworm (WBC) and fall armyworm (FAW). The project has been conducted from Hale to Parmer and up to Dallam and across to Lipscomb by Texas A&M AgriLife County Extension agents. In 2014 eleven county Extension agents monitored moth activity in thirteen High Plains counties. Also, Mr. Haden Hadley, student intern, monitored moth activity in Parmer County. Three pheromone bucket style traps were setup (one per pest species) in each of 27 corn fields. This provided a total of 81 traps across the Texas High Plains

for monitoring the real time abundance and flight duration for each of the three moth species. Each moth species have distinctively different moth flights during the growing season. And, there was considerable variability in a moth's abundance across counties in 2014 and differences from previous years. SWCB activity in 2014 was similar to 2013, but lower compared to 2011 and 2012. Counties with the lightest activity in 2014 were Gray, Hale, Hutchinson, Parmer, Randall, and Swisher. In contrast, Deaf Smith, Dallam, Hansford, Hartley, and Moore counties had much heavier SWCB activity. The SWCB flights of second generation moths occurred from July 22 to August 19. The western bean cutworm flight activity in 2014 occurred from July 1 to August 12, which was again similar to 2013 but much later in the season than 2011 and 2012. WBC moth captures were heaviest in Dallam, Hartley, Hansford, as and Moore counties. WBC were monitored farther east as Lipscomb and Ochiltree counties and south into Castro, Deaf Smith, Gray, Parmer, Randall and Swisher counties. Fall armyworm moths were captured in even higher numbers throughout the growing season in 2014 compared 2013. FAW moth flights were extremely high in Gray, Lipscomb, and Randall counties.

By monitoring the moth activity and reporting the findings through Texas A&M AgriLife Extension newsletters, news articles and phone contacts to producers, crop consultants, local ag suppliers, and ag-aviators, they were able to assess when infestations were a potential threat for making management decisions. From the response of farmers, crop consultants, aerial applicators and other individuals the moth trapping project is meeting a critical need for corn producers in the Texas High Plains.

Introduction

There are approximately 1 million acres of corn grown in the Texas High Plains yearly. Producers that plant non-Bt corn for refuge requirements and for human food consumption are vulnerable to heavy damage from southwestern corn borer (SWCB), western bean cutworm (WBC), and fall armyworm (FAW) infestations. Depending on the Bt-corn hybrid a producer plants, a certain percentage of the corn acreage has to be planted to non-Bt corn hybrids as a refuge to prevent these corn pests from developing resistance to the Bt corn toxins. For corn grown in cotton producing areas (south of Amarillo, TX) the refuge acreage is 20% to 50% non-Bt corn acreage. Fields in non-cotton areas (north of Amarillo, TX) the refuge area is 5% to 20% non-Bt corn acreage or a 95% Bt – 5% non-Bt refuge in a bag hybrid. Also, some of the Bt corn hybrids with single gene toxins do not provide 100% protection against WBC and FAW infestations resulting in corn kernels being damaged from larvae feeding in the ear. Recently, there has been an increased incidence of damage even to the Bt-Herculex corn hybrid. Some food grade corn hybrids do not have the Bt technology and if a producer selects these hybrids then 100% of the corn acreage is susceptible to damage from these pests. Therefore, if just 20% of all corn grown on the Texas High Plains there can be 200,000 acres of corn annually not protected from these corn pests.

Knowing the moth activity during the growing season is critical to making informed management decisions. The activity of these three corn pests can occur at different times and at different infestation levels depending on seasonal weather conditions. This makes it difficult for producers, crop consultants, local ag suppliers, and ag-aviators to know when there will be damaging infestations and when to make timely insecticide applications for optimum control to minimize economic losses. Therefore to assist producers, crop consultants, local ag suppliers, and ag-aviators with knowing when these pests are active, a network of Texas A&M AgriLife

County Extension Agents (CEA) across the Texas High Plains was organized to monitor the moth flight activity of SWCB, WBC, and FAW.

Objective

To provide current real time information to corn producers, crop consultants, local ag suppliers, and ag-aviators throughout the Texas High Plains about the activity of Southwestern corn borer (SWCB), Western bean cutworm (WBC) and fall armyworm (FAW) moth flights during the 2012 growing season.

Method and Materials

Eleven county Extension agents setup pheromone bucket style traps in 13 Texas counties from Hale to Parmer and up to Dallam and across to Lipscomb to monitor the abundance and duration of the moth activity. Also, a student intern setup traps in 2 fields in Parmer county. A total of 81 traps (one per pest species) were setup in 27 corn producer's fields and monitored weekly starting June until the end of August. A spreadsheet with graphs was setup on google documents so each of the county extension agents could post data from their counties. Trap catches from each field in a county were summarized and made available weekly to producers, crop consultants, local ag suppliers, and ag-aviators through phone calls and text messages from the local county extension agents, newspaper articles, county extension agent newsletters, the Texas AgriLife Extension Panhandle Pest Update newsletters, and postings on the Texas AgriLife Extension website Insect Surveys (<http://amarillo.tamu.edu/facultystaff/ed-bynum/insects/>), Texas Panhandle Pest Update blog (<http://txppipm.blogspot.com>) , and on Texas Panhandle IPM twitter (<https://twitter.com/txpipm>).

Results and Discussion

Moth Trapping

Climatic conditions change from year to year which influences the flight activity and abundance of moths. Comparing moth trap catches between 2011 to 2014 show the importance of having a monitoring system each year to determine moth activity of SWCB, WBC, and FAW. The moth activity is really unique to a county and can be variable from year to year. In 2011, SWCB moths increased to extremely high numbers (3,000 to 5,000 per weekly trap catch) in Deaf Smith County and continued for an extended period of time from July 18 to August 29. The extended flight of SWCB moths indicated there was a 3rd generation of SWCB in 2011. In Dallam County, high numbers of SWCB moths were also trapped from July 25 to August 15 (Fig. 1). Comparatively, moderate SWCB activity was recorded in Sherman County while the remaining counties had relatively low levels of SWCB moth activity. In 2012, SWCB moth flights did not reach the high levels that were present in 2011 (Fig 2). The second generation moth activity across the Texas High Plains was basically from July 17 to August 14. Counties with the highest numbers of SWCB moths during this time were Lipscomb, Parmer, and Dallam (Fig. 4). Dallam county had an early peak of SWCB (June 29) and Parmer county had a late flight of SWCB (Aug 28 to Sept 4). Counties with more moderate, but significant, SWCB flights were Deaf Smith, Hansford, Hutchinson, and Sherman. Counties with little or no SWCB activity were Gray, Hale, Ochiltree, Potter, Randal and Swisher. In 2013 and 2014 the overall densities

of SWCB moths were similar but much lower than in 2011 and 2012 (Fig. 3 and Fig. 4). Still larval infestations from moths in both years were high enough to cause economic losses in fields across the High Plains. In both years, counties with the lightest activity were Gray, Hale, Hutchinson, Randall, and Swisher. In contrast, Deaf Smith, Dallam, Hartley, and Moore counties had much heavier activity of SWCB both years. Hansford, Ochiltree, and Lipscomb had moderate levels of moth activity in 2013 and much lower activity in 2014. The flight activity of SWCB moths peaked a week earlier in 2013 compared to 2014.

WBC moth flight activity in 2011 was from June 14 to August 1 with predominately high numbers from June 27 to July 25 in Dallam, Hartley and Sherman counties (Fig. 5). Although moth trap catches were not extremely high in Moore County, trap catches showed WBC moths were present and active during this time. The rest of the counties had nominal to no activity of WBC. In contrast, WBC moth activity in 2012 was predominately for June 19 to July 10, but moths continued to be activity in low densities until August 7 (Fig. 6). This continued activity extended the application window for farmers to protect fields from larval infestations. In 2013, WBC moths captures were heaviest in Dallam, Hartley, Hansford, and Moore counties (Fig.7). The flight activity of western bean cutworm moths did not begin until July 2 but continued until August 13. In 2014, the moth flight began July 1, about the same time as 2013 (Fig. 8), but the number of moths captured peaked earlier and were higher than in 2013. Conditions must have been favorable for WBC movement because moths were even captured in moderate levels in Deaf Smith, Castro, and Swisher counties.

The pheromone lure for FAW moths is not as attractive to moths as lures for SWCB and WBC. Still the lures are effective enough to show patterns of moth flights during the growing season. FAW moth activity in 2011 began with relatively high numbers as shown by trap numbers June 6 in several counties, but activity dropped to low levels the rest of the growing season.(Fig 9). In 2012, FAW activity was relatively low from June 5 to August 21 (Fig. 10). From August 28 to September 18, FAW moth activity increased in during this time, particularly in higher numbers in Deaf Smith and Parmer counties. The trap catches in Deaf Smith and Parmer counties were extended into September at the request of farmers wanting more information about late season activity of FAW. In 2013, FAW moths were captured in high numbers in Gray, Lipscomb, Parmer, and Randall counties throughout the growing season (Fig. 11). The FAW moth activity peaked on July 2 and there was another flight from Aug. 13 to Sept. 3. County Extension Agents from Deaf Smith and Lipscomb counties continued monitoring FAW until September 24 to provide information about late season activity of FAW. In 2014, the FAW moth activity was the heaviest compared to any year that this monitoring project has been capturing moths (Fig. 12). FAW activity was widespread across the Texas High Plains. There were three peaks of moth activity (June 24, July 15, and Aug. 19) during the growing season. Counties with the overall highest number of FAW moths were Lipscomb, Randall, Gray, Parmer, Ochiltree, and Deaf Smith counties.

These moth trapping data demonstrate the variability and differences of flight patterns of the three moth species. Monitoring moth activity reveals that moth activity is different each year area wide and within the counties. As farmers extend their planting dates into June, there will be fields across the Texas High Plains there will be different time periods when fields are vulnerable to egg lay and larval damage from each of these different moths. Therefore, by monitoring the moth activity and reporting the findings to producers, crop consultants, local ag suppliers, and ag-aviators, they are able to assess when infestations are a potential threat and when activity is not a threat.

Impact of Moth Trapping Project

After the 2011 growing season was completed the results of the moth trapping survey project was reported to producers at several meetings sponsored by the county extension agents. These meetings were in Dalhart, Canyon, and Dumas, Stratford, Perryton, Morris, Pampa, and Hereford. These meetings increased individual awareness of the moth trapping project. At the meetings farmers and crop consultants wanted to know if the project was going to be continued in 2013. Their comments were that they liked having the data to help them confirm what they were seeing and to know what was happening in other areas of the Texas High Plains.

In 2012, farmers and consultants in Deaf Smith and Hutchinson request that their county agent increase the number of fields monitored and wanted traps on their fields. After visiting with consultants and farmers, the moth data from the trapping project was spread more by word of mouth. After each Panhandle Pest Update newsletter that contained moth trapping data, Mr. Russell French, Pioneer seed company representative, would forward the newsletter to other farmers, crop consultants, and company representatives on his email list.

In 2013, the response from farmers and consultants has continued to be very positive. Mr. Russell French continued forwarding the Panhandle Pest Update newsletter to his clientele list because of the importance of the moth trapping data to producers. From the response of farmers, crop consultants, aerial applicators and other individuals the moth trapping project is meeting a critical need to corn producers in the Texas High Plains.

In 2014, I setup a blog, Texas Panhandle Pest Update blog (<http://txppipm.blogspot.com>), and a twitter account, Texas Panhandle IPM twitter (<https://twitter.com/txpipm>) to better distribute information about the moth trapping project. I take the information from my Texas Panhandle Pest Update Newsletter and put it in the Texas Panhandle Pest Update blog. Then I tweet a short statement about a particular issue and direct it to my blog. Mr. Kerry Todd, Account Manager for Pioneer Hi-Bred has re-tweeted information to his clientele. I have 51 followers, but Mr. Todd has 540 followers. These new social media venues provide more opportunities for distributing the moth trapping monitoring project. Crop consultants, aerial applicators, and local ag-suppliers continue to request the weekly moth trapping results.

During all four growing seasons the moth trapping data from weekly catches were distributed to farmers, crop consultants, and agri-business individuals on the Panhandle Pest Update newsletter e-mail list. But, County Extension agents are the main source for providing trapping data to their farmers, crop consultants, and agri-business individuals. A survey was conducted after the 2011 season to evaluate the importance of the trapping project to farmers, crop consultants, aerial applicators, and agribusinesses. From the survey, 54.5% of the respondents rated the trapping project as important and 36.4% rated the project as very important for a 90.9% satisfaction rating of the project. There were producers (45.4%), crop consultants (36.4%), local ag suppliers (18.2%), and aerial aviators (27.3%) responding to the questionnaire and as can be seen from those responding some individuals designated more than one occupation. One hundred percent of those responding indicated they got their information from their local County Extension agent by phone calls, text messages and other ways. And, 36.4% got information from The Texas A&M AgriLife Extension Panhandle Pest Update newsletter or from the Texas A&M AgriLife Extension website – Insect Surveys. The trapping data was used to help determine if the moth activity of a particular species (SWCB, WBC, FAW) was a threat or not a threat from one week to the next. This survey showed that individuals used the information in making management decisions. Forty-five percent indicated the moth trapping

data helped them determine if the moth activity was a threat and another 45% indicated the data helped them determine that the moth activity was not a threat. If moths were not a threat 36% of the individuals indicated the information prevented them from making a spray application. But, when moth activity was a potential threat individuals scouted fields more frequently (54.5%), it helped them make better timing of their spray applications (27.3%), and 9% indicated there was a need to increase the number of spray applications based on the moth trapping information. Comments from those filling out the questionnaire were that the moth trapping data “coincided with egg laying”, “it confirms what I was seeing”,

This year, 2014, Mr. Rick Auckerman, CEA – Deaf Smith County, gave out the questionnaire to individuals in his county. Many of the individuals that responded designated they had more than one occupation. Thirty-seven percent marked that they were producers, 62.5% marked crop advisor, 50% marked local ag supplier, and 37.5% marked they were ag aviators. Again, 100% indicated they got their information from Mr. Auckerman either by phone calls, text messages, or email. Twenty-five percent got their information from Mr. Auckerman’s newsletter and another 25% got their information from The Texas A&M AgriLife Extension Panhandle Pest Update newsletter or from the Texas A&M AgriLife Extension website – Insect Surveys. Eighty-eight percent rated the value of the moth trapping data to be very important and 12% rated the data to be important. For the question of whether or not the moth trapping data helped in determining if the moth activity was a threat or not a threat, 87.5% marked that the data helped determine the moth activity was a threat to pending infestations. Although a low percentage of the responders (12.5%) marked moth activity was not a threat, 37.5% indicated the moth trapping data prevented a spray application. When moth activity was a threat individuals marked that they scouted fields more frequently (62.5%). And, 100% indicated the information helped make better timing of spray applications. There was an additional question in this survey that asked individuals to put an economic value (\$/acre) of the moth trapping data to their situation. Most of the responders indicated the value was \$10 to \$20 per acre, but for a few the value of the project was up to \$100 per acre. Comments were “confirmation of field observations”, “gives us an idea of when and where we could have activity on each field”, and “helps me make timely decisions on when to spray”.

These surveys have shown that this moth-trapping project is helping individuals make more informed management decisions for the three moth species being monitored across the Texas High Plains.

Figure 6. Western Bean Cutworm 2012




Figure 7. Western Bean Cutworm - 2013

